

IOWA PRENATAL CARE BARRIERS PROJECT

DATA FROM 2017 SURVEYS

2017 Respondents from All Hospitals in Iowa: 20,919

2017 Respondents with Babies Born in **Woodbury County: 249**

Demographic Indicators	State	County
Mother's Age		
Less than 18 years	01%	<01%
18-19 years	03%	05%
20-25 years	25%	34%
26-30 years	36%	31%
31-35 years	26%	21%
More than 35 years	09%	09%
Mother's Ethnicity		
Hispanic	07%	17%
Non-Hispanic	93%	83%
White	90%	86%
Black	06%	05%
Asian or Pacific Islander	03%	03%
American Indian or Native Alaskan	01%	05%
Socioeconomic Indicators		
Mother's Education		
Less than high school	07%	08%
High school	21%	26%
Some college	34%	37%
College graduate (Bachelor's degree)	27%	19%
Graduate school	12%	09%
Household Income		
Less than \$10,000	12%	20%
\$10,000 - \$19,999	09%	17%
\$20,000 - \$29,999	09%	08%
\$30,000 - \$39,999	08%	07%
\$40,000 - \$49,999	07%	08%
\$50,000 or more	54%	40%

Information Received in 12 Months Prior to Becoming Pregnant	State	County
Health Care Provider Talked About:		
Birth control	49%	43%
Timing of pregnancy	26%	23%
Spacing of children	11%	10%
Doctor, Nurse, or Other Health Care Provider Talked About:		
Taking folic acid or a multivitamin	44%	43%
Smoking	29%	32%
Drinking alcohol	25%	26%
Your weight	24%	24%
Regular exercise	27%	27%
How romantic relationships affect your health	10%	07%
Oral Health	23%	19%
Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant	31%	35%
Behavioral Indicators During Pregnancy	State	County
No Birth Control was Used at Time of Conception	86%	85%
Unintended Pregnancy and Did NOT Use Birth Control	70%	70%
Cigarettes Smoked During Last 3 Months of Pregnancy		
Not smoked in past 2 years	78%	70%
Zero cigarettes per day	53%	52%
1-10 per day	41%	42%
11-20 per day	05%	05%
More than 20 per day	01%	<01%
Health Care Services During Pregnancy	State	County
Visited a Dentist, Dental Clinic, or Received Dental Care	56%	49%
Diagnosed with Depression During Pregnancy	03%	01%
Took Prescription Medication for Depression	48%	42%
Received Counseling for Depression During Pregnancy	30%	35%
HIV/AIDS Testing During Pregnancy		
Yes	41%	44%
No	32%	30%
No, Testing was offered, but did not want to be tested	02%	02%
Don't know	25%	24%

Received Flu Shot During Pregnancy

Yes	65%	72%
No, Received flu shot before became pregnant	07%	04%
No, Did not want flu shot during pregnancy	12%	12%
No, Doctor said to wait until after pregnancy	01%	<01%
No, Do not ever get flu shots	14%	10%
No, Flu shot was not available	02%	02%

Hospital Services**State****County****Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby**

Yes	84%	84%
No	13%	14%
Don't remember	02%	02%

During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss:

Action steps to decrease frustration when infant cries	45%	67%
How to lay the baby down to sleep	58%	68%
Dangers of shaking an infant	43%	66%
Did not receive any information and have not scheduled a time to discuss	30%	16%

Things that may have happened at the hospital where the birth occurred

I breastfed my baby in the hospital	81%	76%
Hospital staff gave me information about breastfeeding	87%	87%
My baby was placed in skin-to-skin contact within the first hour of life	84%	82%
I breastfed in the first hour after my baby was born	64%	63%
Hospital staff helped me learn how to breastfeed	72%	72%
Hospital staff told me to breastfeed whenever my baby wanted	68%	66%
I am feeding my baby	94%	91%

Technical Note: Because percentages were rounded, the sum of values may not equal 100%.

For more information: Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105.

Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2018 Data.